

# Antimicrobial Resistance (AMR): An Ayurvedic Insight

Sijna V.P.<sup>1</sup>, Anjana R.<sup>2</sup>, Archana Radhakrishnan<sup>3</sup>, Vineetha N.<sup>4</sup>, Jaseena C.<sup>5</sup>

<sup>1</sup>Assistant Professor, Department of Swasthavritta

<sup>2</sup>Assistant Professor, Department of Kaumarbhritya

<sup>3</sup>Assistant Professor, Department of Rachana Sharir

<sup>4</sup>Associate Professor, Department of Kayachikitsa

<sup>5</sup>Associate Professor, Department of Kriya Sharir,

KMCT Ayurveda Medical College, Manassery, Kozhikode, Kerala - 673 602

Corresponding Author: Sijna V.P.

## ABSTRACT

Antimicrobial resistance (AMR) is the ability of a microbe to resist the effects of medication that once could successfully treat the microbe. Antimicrobial prophylaxis is commonly used by clinicians for the prevention of numerous infectious, but its use should be limited to specific, well-accepted indications to avoid excess cost, toxicity, and antimicrobial resistance. Rasayana drugs helps in improving immunity as well as physical strength by acting as body nourisher, improves complexion and prevents degenerative changes. Due to the improvement of immunity, microbes cannot colonize the body easily. Hence it will be protective for the body. The paper summarizes the Ayurvedic approaches which can be used effectively for prevention and treatment of bacterial diseases.

**Keywords:** Antimicrobial resistance, AMR, Ayurveda, antimicrobial resistance, immunity

## INTRODUCTION

A microorganism or microbe is a microscopic organism, that exists in its single-celled form or in a colony of cells. The existence of unseen microbes was suspected from ancient times.<sup>1</sup> These single celled creatures have threatened humanity with infection which had high mortality and morbidity.<sup>2</sup> These microorganisms have potential to spread infection at a very rapid rate and to cause widespread epidemics.<sup>3</sup> In 1928, Alexander Fleming discovered the first antibiotic penicillin. By discovering this magic bullet, Fleming set the wheels in motion to create one of the most useful class of drugs in medical history.<sup>4</sup> Next In the next three decades there was development and discovery of a wide variety of antimicrobial agents.<sup>5</sup> In Fleming's Nobel Prize acceptance speech, he warned that the overuse of penicillins might, one day, lead

to bacterial resistance which will be a big problem.<sup>6</sup>

Antimicrobial resistance (AMR) is one of the world's most serious public health problems.<sup>7</sup> Many of the microbes (bacteria, viruses, protozoa) that cause infectious disease no longer respond to common antimicrobial drugs (antibacterial drugs including antibiotics, antiviral and antiprotozoal drugs).<sup>8</sup> Antimicrobial resistance (AMR or AR) is the ability of a microbe to resist the effects of medication that once could successfully treat the microbe.<sup>9</sup> The term antibiotic resistance (ABR) is a subset of AMR, as it applies only to bacteria becoming resistant to antibiotics. Resistant microbes are more difficult to treat, requiring alternative medications or higher doses of antimicrobials. These approaches may be more expensive, more toxic or both. Microbes resistant to multiple

antimicrobials are called multidrug resistant (MDR). Those considered extensively drug resistant (XDR) or totally drug-resistant (TDR) are sometimes called 'superbugs'.

Antimicrobial prophylaxis is commonly used by clinicians for the prevention of numerous infectious, but its use should be limited to specific, well-accepted indications to avoid excess cost, toxicity, and antimicrobial resistance. Antimicrobial prophylaxis may be considered primary (prevention of an initial infection) or secondary (prevention of the recurrence or reactivation of an infection

### Action of Antibiotic

Different antibiotics have different modes of action, owing to the nature of their structure and degree of affinity to certain target sites within bacterial cells.

1. **Inhibitors of cell wall synthesis.** While the cells of humans and animals do not have cell walls, this structure is critical for the life and survival of bacterial species. A drug that targets cell walls can therefore selectively kill or inhibit bacterial organisms. Examples: penicillins, cephalosporins, bacitracin and vancomycin.
2. **Inhibitors of cell membrane function.** Cell membranes are important barriers that segregate and regulate the intra- and extracellular flow of substances. A disruption or damage to this structure could result in leakage of important solutes essential for the cell's survival. Because this structure is found in both eukaryotic and prokaryotic cells, the action of this class of antibiotic are often poorly selective and can often be toxic for systemic use in the mammalian host. Most clinical usage is therefore limited to topical applications. Examples: polymyxin B and colistin.
3. **Inhibitors of protein synthesis.** Enzymes and cellular structures are primarily made of proteins. Protein synthesis is an essential process necessary for the multiplication and

survival of all bacterial cells. Several types of antibacterial agents target bacterial protein synthesis by binding to either the 30S or 50S subunits of the intracellular ribosomes. This activity then results in the disruption of the normal cellular metabolism of the bacteria, and consequently leads to the death of the organism or the inhibition of its growth and multiplication. Examples: Aminoglycosides, macrolides, lincosamides, streptogramins, chloramphenicol, tetracyclines.

4. **Inhibitors of nucleic acid synthesis.** DNA and RNA are keys to the replication of all living forms, including bacteria. Some antibiotics work by binding to components involved in the process of DNA or RNA synthesis, which causes interference of the normal cellular processes which will ultimately compromise bacterial multiplication and survival. Examples: quinolones, metronidazole, and rifampin.
5. **Inhibitors of other metabolic processes.** Other antibiotics act on selected cellular processes essential for the survival of the bacterial pathogens. For example, both sulfonamides and trimethoprim disrupt the folic acid pathway, which is a necessary step for bacteria to produce precursors important for DNA synthesis. Sulfonamides target and bind to dihydropteroate synthase, trimethoprim inhibit dihydrofolate reductase; both of these enzymes are essential for the production of folic acid, a vitamin synthesized by bacteria, but not humans.

### Emergence of antibiotic resistance

India has emerged as the world's largest consumer of antibiotics with a 62 % increase in popping habits over the last decade. The study "Global Trends in Antibiotic Consumption, 2000-2010," by scientists from Princeton University has found that worldwide antibiotic use has risen a staggering 36% over those 10 years, with five countries – Brazil, Russia, India,

China and South Africa (BRICS) – responsible for more than three quarters of that surge.

### **Antimicrobial resistance in India**

Different data collected from various researches shows that bacteria attain resistance power against most of the widely used antibiotic in India. This may cause some serious health challenges including inadequate empiric antibacterial therapy, treatment failure, problems of infection control, increased treatment expenses and increase in the spread of infectious disease caused by resistant microbes.

### **Factors of Antimicrobial Resistance**

#### **Environmental factors**

Over the past years, the role of the environment as an important source of spread of resistance has been increasingly identified.<sup>10</sup> Apart from human transmission, environmental dissemination routes for resistant bacteria have also been pointed out as potential source for the spread. Environmental factors mainly including soil-related factors, animal husbandry and waste management lead to main contributing factors for AMR.<sup>11</sup>

#### **Drug related factors**

On the basis of various researches, human antimicrobial misuse is regarded as the greatest contributing factor toward antimicrobial resistance.<sup>12</sup> This is mainly due to the use of over the counter medicine without any prescription from registered medical practitioners or by irrational fix dose by practitioner. Counterfeit drugs and use of sub-standard drug acts also act as major contributing factors for induction of antimicrobial resistance.

#### **Patient related factors**

Irrational use of antibiotics may increase the resistance in individuals, on the community level and in the society as a whole. Increased trends of self-medication and poor adherence of dosage may also lead to antimicrobial resistance.<sup>13</sup>

### **Impact of antimicrobial resistance**

The World Health Organization (WHO) published the first global surveillance report on antibiotic resistance (ABR) in 2014 to shown that five out of the six WHO regions had more than 50% resistance to third generation cephalosporins and fluoroquinolones in *Escherichia coli* and methicillin resistance in *Staphylococcus aureus* in hospital settings. Similarly, more than 50% resistance to third generation cephalosporins and carbapenems was reported in *Klebsiella pneumoniae*. The report attributed 45% of deaths in both Africa and South-East Asia to multi-drug resistant (MDR) bacteria. It further revealed that *K. pneumoniae* resistant to third generation cephalosporins was associated with elevated deaths in Africa (77%), the Eastern Mediterranean region (50%), South East Asia (81%) and Western Pacific region (72%).<sup>14</sup>

Several resistant bacteria have been increasingly involved in infectious diseases in humans, specifically, *Enterococcus* spp., *S. aureus*, *K. pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacter* spp. They are collectively termed ESKAPE and recently gained further global attention by being listed by the WHO as priority antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics.<sup>15</sup> All these global impact may produce major consequences related to public health in terms of longer hospital stay and increased treatment duration. It also leads to increased overall health care expenditure and reduced productive time resulting in decreased quality of life.

### **Concept of microbiology in classics**

The microorganism and their role in producing various diseases have been elaborately described in Ayurvedic science. There are many references related to microbes in ancient literature. In Rigveda, microorganisms are described as invisible organisms having specific unique characteristics. And in Atharvaveda it is mentioned as a sukshma jantu. The

classical Ayurvedic texts have documented fascinating observations about pathogenic organisms. Charaka samhitha classifies organisms into normal (sahaja) and the pathogenic or invaders (agantu). The pathogenic parasites are further classified into those that grow on the external surface or internal environment of the body termed as krimi. In various Ayurvedic classics, Acharyas have beautifully explained the term graha which capture human and produce variety of symptoms. Along with this Acharyas mentioned the spread of these microbes in terms of Janapadhodhwamsa and Aupasagika vyadhi.

### Handling AMR threat through Ayurveda

Ayurveda promotes conceptual approach to *Svasthya* by stabilizing the delicate balance of nature. Ayurveda approaches act on the root cause of the disease rather than its symptoms. Through these principles, Ayurveda strategies against antimicrobial resistance can be pointed into three major objectives include-

- 1) Decrease the disease transmission
- 2) Reduce the use of antibiotic
- 3) Proper care after antibiotic use

For acquiring these major goals we can divide Ayurveda management into preventive, curative and convalescent care.

### Prevention of antimicrobial resistance

In modern epidemiology, the concept of disease causation is explained in terms of epidemiological triad in which agent, host and environmental factors are interrelated in different complex ways to produce an infectious disease. So the prevention of any disease is mainly through the breakage of this triad. This may be achieved in three important steps:

- Source reduction
- Weakening the mode of transmission
- Strengthening the host

Prevention is the measures taken by an individual prior to the onset of any disease, which will remove the possibility of the disease occurring in the future. Ayurvedic principles related to prevention is

widely accepted globally due to its safety and cost-effectiveness. Significance of modes of intervention in prevention focus on the pre-pathogenic phase. Prevention at this step is mainly categorized into primordial and primary prevention. Primordial prevention is achieved by health education. Modes of intervention in primary prevention can be focused on health promotion and specific protection. The following points mentioned can be considered for the concept of health education in primordial and primary prevention:

- a. Dinacharya - Daily regimen to be followed by an individual.
- b. Ritucharya - Seasonal regimen to be followed by an individual.
- c. Samshodhana - Seasonal cleansing of the individual.
- d. Adhaaraniya Vegas - Non- retention of the Natural urges.
- e. Rasayanas - Intake of Rasayanas.
- f. Sattvavijaya - Improving the mental strength with the help of Yoga and Dhyana

Health promotion concerns activities within as well as outside the health sector which include health education to increase awareness of health problems so that populations identify their health needs and become familiar with preventive strategies and the health facilities available. This is the only component which has a long-term and lasting benefit. For prevention against microbes, promote health education related to Ayurvedic basic principles like dinacharya and Acharya rasayana, Along with health education one should follow Ayurvedic specific protective measures like jala shodhana and vayushodhana.

For strengthening the host, promote health education related to daily regimen, night regimen, seasonal regimens and Acharya rasayana.<sup>16</sup> For the specific protection one should always consumes wholesome food, be aware about the ahara matra and kaala. Charaka emphasizes on food which are wholesome to the body like rice, green gram, rock salt, goose berry,

barley, rain water, milk, ghee, meat and honey. These should be consumed regularly for maintenance of health and prevention of diseases. One should avoid Prajnaparadha (intellectual errors), take care of sense organs, by avoiding under utilization, wrong utilization, over utilization of sense organs, and one should have good memory, knowledge about place, time and one self and should follow rules of Sadvritta to prevent diseases. The diseases due to variations in the climate can be prevented by following purification in respective seasons.<sup>17</sup> The rejuvenation therapy (Rasayana chikitsa) is used in the prevention and cure of disease. Rasayana like aswagandha choorna, chyavanaprasa, indukantha gritha and amalaki rasayana have its own proven immune modulatory effects against the development of infectious diseases. So by consuming all these things one should attain bala against disease and break the epidemiological triad.

### Curative care through Ayurveda

Studies conducted in various countries have correlated antibiotic consumption with the prevalence of antibiotic resistance. Apakarsana, praktivighāta and nidāna-parivarjana are the basic treatment measures mentioned in Ayurveda classics against microbes. Therapeutically it can be correlated the term śodhana, śamana and nidāna-parivarjana respectively.<sup>18</sup> Panchakarma like Śirovirecana, vamana, virecana and āsthāpana are pointed for apakarsana therapy.<sup>19</sup> Antagonist drug therapy for destruction of infectious agent and reduction of symptoms is achieved through praktivighāta and is performed through the drugs used in krimighna and jvarahara like Mahākashāyas.<sup>20</sup> Mahakashayas mentioned in our Ayurvedic classics are having Visaghna, Vranaśodhana, Vranaropana and Kleda-pūyopaśosana activities. The ultimate aim of all these drug is to arrest and encounter the infection. All these antimicrobial drugs are used in the post pathogenesis phase and main action of these

drug are to encounter the visha or antigen caused by the microbes symptoms produced due to the microbes like kleda, pūya, jvara, kandu, dāha etc. Krimighna, Kandughna, Kusthaghna, Jvarahara, Śvāsahara, Kāsahara, Śothahara, Śītāpraśamana Mahākashāyas are to be used against these symptoms.<sup>21</sup> Some groups of drugs in Mahakashayas are also used to arrest the infections caused by specific type of microbes which produce different types of discharges, burning sensations, pain, redness etc. Āragvadhādi mahakashaya having Kushtagna and Kandugna activity destroy the microbes, alleviates itching and cleanses wound; Sālasārādi group is also having Kushtagna activity and administered in various types of infective skin diseases; Varunadi group is having Vranaghna activity and is highly effective in the treatment of internal abscess; Lodhrādi are yonidoshahara group which arrests the diseases of female genital tract caused by different pathogens; arkādi group help to cleanses infective wound; Surasādi group is highly effective in respiratory infection both upper and lower tract; Pippalādi group is effective in acute and chronic rhinitis; Elādi group is highly effective in boils and furuncles; Vacādi and Haridrādi both encounter the pathogens in the diseases like diarrhoeal disorders; Parusakādi is advocated in urinary disorders; Priyaggvādi and Ambasthādi both are useful in chronic type of dysentery and also effective in wound healing; Nyagrodhādi groups are beneficial for chronic wound arrests the infections in female genital tract; Mustādi group has positive result in female genital tract infections too; Lāksādi group is useful in infective wound and act as anti-helminthic; Trinapachamūla is highly effective in urinary tract infections. All the above conditions and or diseases which are arrested through the administration of different types of said groups of drugs are clinically caused due to infections.<sup>22</sup>

Different Ayurvedic preparations like Mahasudharshana choorna, Triphala choorna, Hareethaki choorna, Ajamodadi



choorna and Dasamoola choorna are having antimicrobial activity against enteric bacterial pathogens. Studies have shown that drugs like Bilva, shalmali, Dadima, Kutaja, Arjuna and Triphala shows strong antibacterial activity against MDR *Salmonella* Typhi.

#### Convalescent care after antibacterial use

Antibiotics can be a very powerful factor causing imbalance of the intestinal microbiota.<sup>23</sup> In 1954, Bohnhoff and co-workers in their studies noticed that mice that were given streptomycin were easily infected by *Salmonella enterica* serovar Enteritidis and introduced the concept that intestinal microbiota could suppress the growth of bacteria that invade mice from the outside through colonization resistance.<sup>24</sup> Direct interaction of intestinal microbiota with bacteria and competition for intestinal nutrients are direct methods of inhibiting the intestinal colonization of pathogens. However, dosing with antibiotics reduces the diversity and abundance of intestinal microbiota, leading to a reduction in the competitive exclusion ability.<sup>25</sup> Indirectly, this destroys the community structure, thereby disturbing the interactions among microbial species and the complementary systems of nutrient metabolic pathways, resulting in widespread fluctuations in the intestinal environment. These changes are not fully reversed, even after several months of discontinuation of dosing.<sup>26</sup> Antibiotic-induced perturbations of the intestinal microbiota alter host susceptibility to enteric infection. Eventually, the antibiotic-induced dysbiosis of the intestinal microbiota affects the development and regulation of the immune system and increases the risk of intestine-related diseases, such as inflammatory bowel diseases and infectious diseases, in addition to diverse immunity-related disorders, such as allergic or atopic skin diseases and type 1 diabetes.<sup>27</sup>

These pathologies in human body may lead to Agnimandhya, Rasa kshaya, Tridosha dushti and finally end in

ojakshaya. So the intervention are mainly focused to three steps

- 1) Repair what has been damaged
- 2) Re introduce beneficial bacteria
- 3) Rejuvenation

After attaining proper agni, next focus is to reintroduce beneficial bacteria into gut flora. For that one should incorporate probiotic in food. Kheera Ghritha Navaneetha and Takra are naturally available probiotics.

#### Rasayana

Rasayana helps in improving immunity as well as physical strength. It acts as body nourisher, improves complexion and prevents degenerative changes. Due to the improvement of immunity, microbes cannot colonize the body easily. Hence it will be protective for the body.

#### CONCLUSION

To conclude, Ayurvedic interventions can be used effectively for prevention and treatment of bacterial diseases. It can also be used in convalescent care subsequent to antibacterial therapy. Hence in the current era of AMR, Ayurvedic interventions will go a long way in reducing the risk of development of AMR and also in alleviating the adverse effects of antibacterial therapy.

#### REFERENCES

1. Karsten Zengler. Central Role of the Cell in Microbial Ecology. *Microbiol Mol Biol Rev.* 2009 Dec; 73(4): 712–729.
2. Johanna F. Lindahl & Delia Grace (2015) The consequences of human actions on risks for infectious diseases: a review, *Infection Ecology & Epidemiology*, 5:1.
3. Thomas V Inglesby and Amesh A. Adalja. Characteristics of Microbes Most Likely to Cause Pandemics and Global Catastrophes. *Global Catastrophic Biological Risks.* 2019; 424: 1–20.
4. Gaynes R. The Discovery of Penicillin- New Insights After More Than 75 Years of Clinical Use. *Emerg Infect Dis.* 2017;23(5): 849-853.

5. Jayachandran S. Pre-antibiotics era to post-antibiotic era. *J Indian Acad Oral Med Radiol* 2018;30:100-1.
6. Noah Rosenblatt-Farrell. The Landscape of Antibiotic Resistance. *Environ Health Perspect.* 2009 Jun; 117(6): A244–A250.
7. Alex Broom, Katherine Kenny, Barbara Prainsack, Jennifer Broom. (2020) Antimicrobial resistance as a problem of values? Views from three continents. *Critical Public Health* 0:0, pages 1-13.
8. Pauline Hood, Ehsan Khan. Antimicrobials. *Understanding Pharmacology in Nursing Practice* . 2019 Sep 4 : 147–165.
9. Aslam B, Wang W, Arshad MI, Khurshid M, Muzammil S, Rasool MH, Nisar MA, Alvi RF, Aslam MA, Qamar MU, Salamat MKF, Baloch Z. Antibiotic resistance: a rundown of a global crisis. *Infect Drug Resist.* 2018;11:1645-1658.
10. Bengtsson-Palme, Johan et al. “ J, Kristiansson E, Larsson DGJ. Environmental factors influencing the development and spread of antibiotic resistance.” *FEMS microbiology reviews* vol. *Microbiol Rev.* 2018;42,1 (2018).(1):fux053. doi:10.1093/femsre/fux053
11. Fletcher S. Understanding the contribution of environmental factors in the spread of antimicrobial resistance. *Environ Health Prev Med.* 2015;20(4):243-252. doi:10.1007/s12199-015-0468-0
12. Castro-Sánchez E, Moore LS, Husson F, Holmes AH. What are the factors driving antimicrobial resistance? Perspectives from a public event in London, England. *BMC Infect Dis.* 2016;16(1):465. Published 2016 Sep 2. doi:10.1186/s12879-016-1810-x
13. Kolozsvári, L.R., Kónya, J., Paget, J. et al. Patient-related factors, antibiotic prescribing and antimicrobial resistance of the commensal *Staphylococcus aureus* and *Streptococcus pneumoniae* in a healthy population - Hungarian results of the APRES study. *BMC Infect Dis* 19, 253 (2019).
14. Founou RC, Founou LL, Essack SY. Clinical and economic impact of antibiotic resistance in developing countries: A systematic review and meta-analysis. *PLoS One.* 2017;12(12):e0189621. Published 2017 Dec 21. doi:10.1371/journal.pone.0189621
15. Kolozsvári LR, Kónya J, Paget J, et al. Patient-related factors, antibiotic prescribing and antimicrobial resistance of the commensal *Staphylococcus aureus* and *Streptococcus pneumoniae* in a healthy population - Hungarian results of the APRES study. *BMC Infect Dis.* 2019; 19(1):253. Published 2019 Mar 13. doi:10.1186/s12879-019-3889-3
16. Vagbhata, Ashtanga Hridaya, Varanasi, Chaukhamba orientalia, 9th edition, 2005, sutra sthana, chapter 4, Sloka 36 pg no-53.
17. Vagbhata, Ashtanga Hridaya, Varanasi, Chaukhamba orientalia, 9th edition, 2005, sutra sthana, chapter 4, Sloka 35 pg no-59.
18. Agnivesh, Vimanasthan, 7th Chapter, 28-30 Shloka, Caraka Samhita, Ayurveda Depika Commentry by Cakrapanidutta, Pt. Yadavji Trikamji Acharya editor, Rastriya Sanskrit Samsasthan , Reprint 2006 Page 26).
19. Agnivesh, Vimanasthan, 7th Chapter, 15th Shloka, Caraka Samhita, Ayurveda Depika Commentry by Cakrapanidutta, Pt. Yadavji Trikamji Acharya editor, Rastriya Sanskrit Samsasthan , Reprint 2006 Page 258-259.
20. Agnivesh, Sutrasthan, 4th Chapter, 11th & 16th Shloka , Caraka Samhita, Ayurveda Depika Commentry by Cakrapanidutta, Pt. Yadavji Trikamji Acharya editor, Rastriya Sanskrit Samsasthan , Reprint 2006 Page 33 & 34 .
21. Agnivesh, Sutrasthan, 4th Chapter, 11-17th Shloka , Caraka Samhita, Ayurveda Depika Commentry by Cakrapanidutta, Pt. Yadavji Trikamji Acharya editor, Rastriya Sanskrit Samsasthan , Reprint 2006 Page 33-34.
22. Sushruta, Sutrasthan , 38th Chapter, 9- 28th shloka, Sushruta Samhita, Nibandhasamgraha commentary by Dalhana, Yadavji Trikamji Acharya editor, Chaukhamba Orientalia, 6th edition, 1997 page 165.
23. Perez-Lopez A, Behnsen J, Nuccio SP, Raffatellu M. Mucosal immunity to pathogenic intestinal bacteria. *Nat Rev Immunol.* 2016;16:135–148.
24. Bohnhoff M, Drake BL, Miller CP. Effect of streptomycin on susceptibility of intestinal tract to experimental *Salmonella* infection. *Proc Soc Exp Biol Med.* 1954;86:132–137.
25. Schubert AM, Sinani H, Schloss PD. Antibiotic-induced alterations of the murine gut microbiota and subsequent effects on

- colonization resistance against *Clostridium difficile*. *MBio*. 2015;6:e00974
26. Sekirov I, Tam NM, Jogova M, Robertson ML, Li Y, Lupp C, et al. Antibiotic-induced perturbations of the intestinal microbiota alter host susceptibility to enteric infection. *Infect Immun*. 2008;76:4726–4736.
27. Yoon, Mi Young, and Sang Sun Yoon. “Disruption of the Gut Ecosystem by Antibiotics.” *Yonsei medical journal* vol. 59,1 (2018): 4-12.
- How to cite this article: Sijna V.P., Anjana R., Radhakrishnan A et.al. Antimicrobial resistance (AMR): an ayurvedic insight. *Int J Health Sci Res*. 2021; 11(1):80-87.

\*\*\*\*\*