



Original Research Article

Albumin/Creatinine Ratio, As Predictor of Microalbuminuria, a Risk Factor For Nephropathy In Type 2 Diabetes Mellitus Patients.

Abdulrahman A Momin¹, Pankaja S Naik¹, Gouri M Bhoite²

¹Department of Biochemistry, Dr. Vasantrao Pawar Medical College, Adgaon, Nashik.

²Department of Biochemistry, Bharati Vidyapeeth University, Dental College, Pune.

*Correspondence Email: rahaman.momin@gmail.com

Received: 14/12/2011

Revised: 24/12/2011

Accepted: 30/12/2011

ABSTRACT

Diabetic nephropathy is characterized by proteinuria and is the leading cause of end-stage renal disease worldwide. A case control study was designed to determine the prevalence of microalbuminuria in type 2 diabetic subjects. 50 subjects with type 2 diabetes and 50 normal healthy controls were screened for albumin and creatinine. The urinary albumin was calculated in terms of ratio with respect to urinary creatinine and expressed as albumin/creatinine ratio (mg/g). Albumin/Creatinine ratio in controls was found to be 11.05 ± 4.52 mg/g. The mean \pm SD of Albumin/Creatinine ratio in total 50 patients was 38.77 ± 23.13 mg/g. The value of urinary albumin between 30 and 300 mg/g of creatinine was considered to be positive for microalbuminuria. Out of total 50 diabetic subjects, 31 patients were found to have albumin excretion of more than 30 mg/g of creatinine in random morning samples and therefore positive for microalbuminuria. The prevalence of microalbuminuria in type 2 diabetic subjects in this study was 62%. Use of the albumin-to-creatinine ratio in an untimed urinary sample should be recommended as the preferred screening strategy for all diabetic patients.

Key words: Microalbuminuria, albumin, creatinine, urine albumin creatinine ratio, type 2 diabetes.

INTRODUCTION

Microalbuminuria is a marker of an increased risk of diabetic nephropathy in patients with Type 1 as well as with Type 2 diabetes. It is common in Type 2 diabetic

patients, especially those of non-European origin. Poor long-term control, hypertension, dyslipidemia and cigarette smoking facilitate the development of diabetic nephropathy.⁽¹⁾ Albumin, a protein having molecular weight of 50,000 is not easily filtered and is not

excreted into urine. This makes albumin excretion into the urine a useful indicator of early glomerular disease.⁽²⁾

Increase in urine albumin seen with diabetic nephropathy can be attributed to degradation of the glomerular basement membranes & hypertension both characteristics of Diabetic Nephropathy.⁽³⁾ Use of the albumin-to-creatinine ratio in an untimed urinary sample is now recommended as the preferred screening strategy for all diabetic patients.^(4,5,6) The effect of volume can be avoided entirely by calculation of the albumin-to-creatinine ratio in an untimed urine specimen. A ratio above 30 mg/g (or 0.03 mg/mg) suggests that albumin excretion is above 30 mg/day and therefore that microalbuminuria is probably present.⁽⁷⁾

Microalbumin has been reported in several studies to predict development of diabetic nephropathy and its mortality risk in diabetic patients. Early detection and aggressive intervention should be given to retard the progression of diabetic nephropathy to end stage renal failure.⁽⁸⁾

MATERIALS AND METHODS

SUBJECTS

The case control study was performed at Department of Biochemistry in Central Clinical Laboratory of MVP's Medical College, Adgaon, Nashik. The study was approved by institutional ethical committee. Total 50 patient samples between age group 40-60 years with Diabetes mellitus, diagnosed according to WHO criteria, from department of medicine, MVP's medical college, Adgaon were collected. The patients were without any history of cardiovascular complications, hypertension or renal diseases, and written informed consent was taken. The study

group was compared with group of age and sex matched 50 control samples. The routine samples in the central clinical laboratory were taken as control samples, the control samples were without any history of diabetes mellitus, nephropathy, cardiovascular diseases and hypertension.

METHODS

The early morning urine samples were collected for the determination of albumin & creatinine and sample for blood glucose in fluoride bulb with potassium oxalate as an anticoagulant. Each urine sample was assayed for creatinine and albumin. The albumin was measured by latex turbidimetry method.^(9, 10) Urine creatinine was measured by Jaffe's kinetic method^(11, 12) and blood glucose by enzymatic GOD-POD method.⁽¹³⁾

RESULTS

The age, blood glucose level, urine albumin, urine creatinine and urine albumin/creatinine ratio were calculated and expressed as mean \pm SD, the results are depicted in Table: 1. The mean age of the patients was 52.6 ± 7.92 years and to that of healthy controls 47.21 ± 9.45 years. The blood glucose level in study group was 298 ± 47.05 mg/dl and highly significant (<0.01) increase is observed as compared to healthy controls i.e. 107 ± 27.39 mg/dl. The urinary albumin level was 62.80 ± 5.31 mg/l in patients and 15.24 ± 0.85 mg/l in controls. Compared to controls the increase in the albumin level in patients was found to be highly significant (<0.01). The difference in urine creatinine level was not found to be significant (>0.05) between patient group (1.62 ± 0.23 g/l) and healthy controls (1.38 ± 0.19 g/l).

Table: 1- The levels of biochemical parameters in study and control group.

Parameter	Patients	Controls	'p' Value
Blood glucose (mg/dl)	298 ± 47.05	107 ± 27.39	P<0.01
Urine albumin (mg/l)	62.80 ± 5.31	15.24 ± 0.85	P<0.01
Urine creatinine (g/l)	1.62 ± 0.23	1.38 ± 0.19	p>0.05
Albumin/creatinine ratio (mg/g)	38.77 ± 23.13	11.05 ± 4.52	P<0.01

The urinary albumin was calculated in terms of ratio with respect to urinary creatinine and expressed as albumin/creatinine ratio (mg/g). Albumin/Creatinine ratio in patients was 38.77 ± 23.13 mg/g, and to that of in controls was 11.05 ± 4.52 mg/g, and this increase in the ratio was found to be highly significant (p<0.01). The ratio of urinary albumin to creatinine between 30 and 300 mg/g is considered to be positive for microalbuminuria. 31 patients were found to have albumin excretion of >30 mg/g of creatinine in random morning samples and therefore positive for microalbuminuria, and confirmed by measurement of microalbumin in 24 hour urine sample.

DISCUSSION

Prevalence of nephropathy in type 2 Diabetes mellitus varies in different population groups i.e., fairly low incidence in Caucasians and a very high incidence in Pima Indians.⁽¹⁴⁾ Microalbuminuria is a marker for diabetic nephropathy in type 2 diabetes mellitus.^(2, 15) Using a specific assay for albumin is a more sensitive technique to confirm nephropathy. Persistent albumin excretion between 30 and 300 mg/day is called microalbuminuria. Fever, exercise, heart failure, hypertension, obesity and poor glycemic control are among the factors that can cause transient microalbuminuria.⁽⁵⁾

Although the 24-hour urine collection was previously the gold standard for the detection of microalbuminuria, it has been suggested that screening can be more simply achieved by a timed urine collection or an early morning specimen to minimize changes in urine volume that occur during the day.^(4, 16)

A value above 30 mg of albumin /gm of creatinine (or 0.03 mg/mg) suggests that albumin excretion is above 30 mg/day and therefore that microalbuminuria is probably present.⁽⁷⁾ In one report by Gross JL⁽⁵⁾ 24-hour urine collections and random, single-void urine specimens for albumin and creatinine were obtained in 14 normal subjects, 13 with type 1 diabetes, and 12 with type 2 diabetes. A close correlation was noted between the two measurements with small variability within patient.

Use of the albumin/creatinine ratio in an untimed urinary sample is now recommended as the preferred screening strategy for all diabetic patients.^(4, 5, 6) Earlier reports have shown good correlation between urinary albumin/creatinine ratio and 24-h urinary excretion in screening for microalbuminuria^(17, 18) and guidelines^(19, 20) suggest that using albumin/creatinine ratio is valid for screening in the diabetic patient.

In the present study the blood sugar level was higher in the study group than healthy controls which was found to be

significant ($p < 0.01$). Urine albumin level in patients was significantly ($p < 0.01$) increased as compared to healthy controls, where as difference of urine creatinine level between study group and healthy controls was not significant. Urine albumin/creatinine ratio was increased in patients as compared to controls, and it is found highly significant ($p < 0.01$).

In a study by Zelmanovitz T,⁽¹⁶⁾ 95 patients with type 2 diabetes found an equally high correlation between 24-hour urine albumin excretion and the albumin/creatinine ratio in the first morning urine.⁽²¹⁾ Mattix⁽²²⁾ et al used albumin/creatinine ratio to calculate microalbuminuria among men and women, the prevalence of microalbuminuria was significantly lower among the men compared with women Waugh⁽²³⁾ et al studied urinary microalbumin/creatinine ratios reference range in uncomplicated pregnancy. Significant increase ($p < 0.016$) in the albumin/creatinine ratio in the third trimester was found by them.

In a cohort study conducted by Cirillo⁽²⁴⁾ et al consisted of 1623 men and women, ages between 45 & 64. They found high albumin/creatinine ratio prevalence, it was 8.5%. The effectiveness of albumin/creatinine ratio was studied by Tazeen⁽²⁵⁾ et al they found, the sensitivity was 46 and specificity was 95%, in women and 60 and 97%, respectively, in men.

This test has the following advantages: it does not require early morning or timed collections, it gives a quantitative result that correlates with the 24-hour urine values over a wide range of protein excretion, it is cheap to perform, and repeat values can be easily obtained to ascertain that microalbuminuria, if present, is persistent.

REFERENCES

1. Casper GS, Nicole L, Harry T, et al.: Amadori Albumin in Type 1 Diabetic Patients Correlation With Markers of Endothelial Function, Association with Diabetic Nephropathy, and Localization in Retinal Capillaries, Diabetes, December 1999;48 (12):2446-2453
2. American diabetes association, Nephropathy in Diabetes, diabetes care, january 2004; 27(1):S79-S83
3. Ghazalli R: Diabetic Nephropathy in Clinical practice guidelines, July 2004: 1-38
4. Mogensen CE, Vestbo E, Poulsen PL, et al.: Microalbuminuria and potential confounders. A review and some observations on variability of urinary albumin excretion. Diabetes Care 1995; 18:572-581
5. Gross JL, de Azevedo MJ, Silveiro SP, et al.: Diabetic nephropathy: diagnosis, prevention, and treatment. Diabetes Care 2005; 28:164-176
6. K/DOQI Clinical Practice Guidelines and Clinical Practice recommendations for diabetes and chronic kidney disease. Am J Kidney Dis 2007; 49(2):S21-S95
7. Nakamura Y, Myers BD: Charge selectivity of proteinuria in diabetic glomerulopathy. Diabetes 1988;37:1202-1211
8. Graziella B, Franco M: Progression to overt nephropathy in type-2 diabetes. The Casale Monferrato study. Diabetes Care 2003; 26:2150-2155
9. Bernard A, Lauwerys R.: Latex immunoassay of urinary albumin. J Clin Chem Clin Biochem, 1983; 21(1):25-30.
10. Burtis CA, Ashwood ER: Tietz textbook of Clinical Chemistry, 3rd edition, WB Saunders Co., 1999.

11. Slot C: Plasma creatinine determination. A new and specific Jaffe reaction method. *Scand J. Clin. Lab Invest*, 1965; 17: 381 – 387.
12. Seaton B, Ali A. Simplified manual high performance clinical chemistry methods for developing countries. *Med Lab Sci*. 1984; 41(4):327–336.
13. Basak A: Development of a rapid and inexpensive plasma glucose estimation by two-point kinetic method based on glucose oxidase-peroxidase enzymes, *Indian Journal of Clinical Biochemistry*, 2007;22 (1): 156-160.
14. Cowie CC, Port FK, Wolfe RA, et al.: Disparities in incidence of diabetic end-stage renal disease according to race and type of diabetes. *N Engl J Med* 1989; 321:1074-1079
15. Gaede P, Tarnow L, Vedel P et al: Remission to normoalbuminuria during multifactorial treatment preserves kidney function in patients with type 2 diabetes and microalbuminuria. *Nephrol Dial Transplant* 2004; 19:2784 –2788.
16. Zelmanovitz T, Gross JL, and Oliveira JR, et al.: The receiver operating characteristics curve in the evaluation of a random urine specimen as a screening test for diabetic nephropathy. *Diabetes Care* 1997; 20:516-519
17. Skov J J, Clausen P, Borch J K et al: Detecting microalbuminuria by urinary albumin/ creatinin concentration ratio. *Nephrol Dial Transplant* 1997; 2:6 –9.
18. Eshøj O, Feldt R B, Larsen M L et al: Comparison of overnight, morning and 24-hour urine collections in the assessment of diabetic microalbuminuria. *Diabet Med* 1987;4:531–533
19. National Kidney Foundation: Clinical practice guidelines for chronic kidney disease: evaluation, classification and stratification. *Am J Kidney Dis* 39 2002; 1:S1– S266
20. American Diabetes Association: Nephropathy in diabetes (Position Statement). *Diabetes Care*, 2004; 27 (1):S79– S83,
21. Fliser D, Wagner KK, Loos A et al.: Chronic angiotensin II receptor blockade reduces (intra)renal vascular resistance in patients with type 2 diabetes. *J Am Soc Nephrol* 2005; 16:1135-1140
22. Mattix HJ, Hsu CY, Shaykevich S et al: Use of the Albumin/Creatinine Ratio to Detect Microalbuminuria: Implications of Sex and Race, *J Am Soc Nephrol* 2002;13: 1034–1039
23. Waugh J, Bell SC, Kilby MD et al: Urinary microalbumin/creatinine ratios: reference range in uncomplicated pregnancy *Clinical Science*, 2003;104: 103–107
24. Cirillo M, Laurenzi M, Mancini M et al: Low Muscular Mass and Overestimation of Microalbuminuria by Urinary Albumin/Creatinine Ratio, *Hypertension*, 2006; 47:56-61
25. Tazeen HJ, Chaturvedi N, Hatcher J et al: Use of albumin creatinine ratio and urine albumin concentration as a screening test for albuminuria in an Indo-Asian population. *Nephrol Dial Transplant*, 2007; 22: 2194–2200
