



Original Research Article

Total and Differential Leucocyte Count in Pregnancy Induced Hypertension

Geraldine Monteiro^{*}, Subbalakshmi NK^{**@}, Anupama N^{**}, Sheila R Pai^{***}

^{*}Tutor and Postgraduate Student, ^{**}Associate Professor, ^{***}Professor,
Department of Physiology, Kasturba Medical College, Manipal University, Mangalore, Karnataka, India.

@Correspondence Email: rao.subbalakshmink@rediffmail.com

Received: 11/10/2013

Revised: 11/11/2013

Accepted: 16/11/2013

ABSTRACT

Introduction and objective: Preeclampsia evokes inflammatory response. But whether inflammatory response observed is in relation to severity of preeclampsia is not clear. This study compared the total and differential leucocyte count between normotensive mothers and mothers with mild and moderate - severe preeclampsia

Materials and methods: Total leucocyte count and differential leucocyte count was done in 92 normotensive mothers, 54 mothers with mild preeclampsia and 28 mothers with moderate-severe preeclampsia.

Results: Mothers with moderate to severe preeclampsia had higher total leucocyte count, differential neutrophil count and differential basophil count compared to normotensive mothers ($P < 0.05$). There was no significant difference in total and differential leucocyte count of mild preeclampsia group compared to normotensive group.

Conclusion: Higher total leucocyte count, differential neutrophil count and basophil count occurs mainly in moderate to severe preeclampsia

Key words: preeclampsia, total leucocyte count, neutrophils, basophils

INTRODUCTION

The hypertensive disorders during pregnancy affect up to 8.0% of all pregnancies⁽¹⁾ and remain a major cause of maternal and neonatal mortality and morbidity worldwide.^(2,3) In women with pre-eclampsia, a history of symptoms is obtained antenatally, with the key alarming clinical symptoms being headache, vomiting, visual disturbances and epigastric

pain. However the unpredictable onset and progression of these symptoms warrants the need for an early and appropriate detection.⁽⁴⁾

Over the years, a lot of interest has been directed at studies on the role of serum uric acid, urea and creatinine in the pathogenesis of pregnancy induced hypertension.^(5,6) However there is a constant ongoing search for better predictors

and prognostic factors to assess the progress and severity of this disease, hematological parameters being one among them.⁽⁷⁾ Additionally pregnancy induced hypertension is a multisystem disorder and may involve inflammatory response.⁽⁸⁾ Therefore this study was undertaken to assess the total leucocyte and differential leucocyte count in mothers with preeclampsia.

MATERIALS AND METHODS

This study was done in teaching hospitals attached to Kasturba Medical College after obtaining permission from ethical committee of the institution overseeing human studies and consent from the study participants.

174 consecutive mothers in 20th week of gestational period were enrolled into the study, in whom fresh medical report on serum protein, total leucocyte and differential leucocyte count was available. Among them 82 were with clinical diagnosis of preeclampsia and 92 were normotensive mothers. Mothers with Pre-eclampsia were further classified into mild (n = 54) and moderate-severe (n =28) based on the blood pressure recording. A blood pressure recording of 140mmHg systolic blood pressure and 90 mm Hg diastolic blood pressure was taken as the cut off point for mild pre-eclampsia while a blood pressure recording of 160 mmHg systolic blood pressure or 100 mm Hg diastolic blood pressure was the starting point of moderate-severe preeclampsia. In this study the diagnosis of preeclampsia was based on the definition of American College of Obstetrics and Gynecologists.⁽⁹⁾

In all the subjects, height and weight was measured and body mass index was calculated using the formula weight in

kilogram divided by height in meters squared. Systolic and diastolic blood pressure was measured in sitting position.

Statistical analysis included analysis of one way variance, Kurskal- Wallis test followed by Dunn's multiple comparison test.

RESULT

Data is presented as mean \pm SD. Data on comparison of baseline characteristics of mothers with preeclampsia and normotensive mothers are presented in table 1. Data on total leucocyte and differential leucocyte count in mild preeclampsia, moderate-severe preeclampsia and normotensive mothers is presented in table 2. Mean age of mothers with mild preeclampsia and moderate-severe preeclampsia was higher compared to normotensive mothers ($p < 0.05$, table 1). Mean age of mothers with mild preeclampsia did not differ significantly compared to mothers with moderate –severe preeclampsia (table 1). Systolic and diastolic blood pressure of mothers with preeclampsia was significantly higher compared to normotensive mothers ($p < 0.001$, table 1). Systolic and diastolic blood pressure of moderate-severe preeclampsia was significantly higher compared to mothers with mild preeclampsia ($p < 0.05$, table 1). Urine protein was significantly higher in mothers with mild and moderate-severe preeclampsia compared to normotensive mothers ($p < 0.001$, table 1). Mean Urine protein of mothers with mild preeclampsia did not differ significantly compared to moderate –severe preeclampsia (table1). Mean body mass index of normotensive mothers and mothers with preeclampsia did not differ significantly (table 1).

Table 1 Baseline characteristics of preeclamptic and normotensive mothers (Values are mean \pm SD).

Variables	Mothers With Mild preeclampsia (n= 54)	Mothers With Moderate – severe preeclampsia (n = 28)	Normotensive mothers (n = 92)	F/KW value	P value
Age (years)	29.44 \pm 3.65*	30.32 \pm 4.11*	27.60 \pm 4.10	11.704	0.0029
Body Mass index (kg/m2)	26.22 \pm 2.93	25.82 \pm 2.70	25.56 \pm 2.65	0.982	0.376
Systolic blood pressure (mmHg)	154.07 \pm 10.00 **†	165 \pm 8.38 **	117.71 \pm 7.12	132.75	< 0.0001
Diastolic blood pressure(mmHg)	95.55 \pm 5.01 **†	105.35 \pm 5.07 **	72.82 \pm 4.52	147.28	< 0.0001
Urine protein	7.17 \pm 1.61 ††	6.21 \pm 1.12 ‡	5.31 \pm 1.07	48.004	< 0.0001

* P< 0.05 compared to normotensive group (Dunn’s multiple comparison).

** p<0.001 compared to normotensive group

Total leucocyte count, differential neutrophil count and differential basophil count of mothers with moderate - severe preeclampsia was significantly higher compared to normotensive mothers (p< 0.05,table 2). Total leucocyte count, differential neutrophil count and differential basophil count of mothers with mild preeclampsia was not significantly different compared to normotensive mothers (table 2). Differential eosinophil count, differential monocyte count and differential lymphocyte count of mothers with preeclampsia did not differ significantly compared to normotensive mothers (table 2).

Table 2. Total and differential leucocyte counts in preeclampsia and normotensive mothers (values are mean \pm SD).

Variables	Mild preeclampsia (n= 54)	Severe preeclampsia (n = 28)	Normotensive (n = 92)	F/kw value	P value
Total leucocyte count	12774.25 \pm 4772.14	13241.42 \pm 4683.81*	10889.67 \pm 3122.78	8.307	0.015
Differential neutrophil count	77.29 \pm 7.74	78.53 \pm 11.418*	75.11 \pm 6.76	6.479	0.039
Differential eosinophil count	0.88 \pm 0.81	0.75 \pm 0.84	1.02 \pm 0.94	2.062	0.356
Differential basophil count	0.55 \pm 0.50	0.82 \pm 0.66*	0.45 \pm 0.52	7.355	0.025
Differential monocyte count	0.88 \pm 0.94	0.71 \pm 0.46	0.47 \pm 0.50	5.293	0.07
Differential lymphocyte count	19.44 \pm 8.38	18.82 \pm 6.48	19.15 \pm 7.31	0.084	0.95

* p< 0.05 compared to normotensive group (Dunn’s multiple comparison).

DISCUSSION

In the present study, urine protein was significantly higher in mothers with preeclampsia compared to normotensive mothers (table 1). But urine protein did not differ significantly between mothers with mild preeclampsia compared to mothers with moderate-severe preeclampsia. This finding suggests that although raised protein is a characteristic feature of pregnancy induced hypertension, this parameter may not be involved in the progression of pathogenesis leading to severe preeclampsia.

Neutrophil count was higher in mothers with moderate to severe preeclampsia compared to normotensive mothers (table 2). Canzoneri BJ et al too have observed increase in total leucocyte count in pregnancy induced hypertensive mothers.⁽¹⁰⁾ Pregnancy induced hypertension

is a multisystem disorder and may involve inflammatory response.⁽⁸⁾ Thus higher leucocyte count especially of neutrophils observed in our study in moderate-severe preeclampsia group could be one of the clinical manifestations of preeclampsia in moderate to severe state but not in mild state of preeclampsia. However, Sivakumar S et al have not observed any difference in total leucocyte count and differential neutrophil count in pregnancy induced hypertensive mothers compared to normotensive mothers.⁽¹¹⁾

In the present study basophil count was significantly higher in mothers with moderate to severe preeclampsia compared to normotensive mothers. Basophils appear in many specific kinds of inflammatory reactions. Thus increased basophil count observed in moderate to severe preeclampsia

may be one of the characteristic features of inflammatory response induced by moderate to severe preeclampsia. However all the studies do not confirm this observation. Lurie S et al did not find any increase in basophil count in mothers with preeclampsia compared to normotensive mothers.⁽¹²⁾ On the other hand Vonne J et al observed decreased basophil count in mothers with preeclampsia compared to normotensive mothers.⁽¹³⁾

CONCLUSION

Our study concludes that total leucocyte count is increased in moderate to severe preeclampsia. Among the differential leucocyte count, neutrophils and basophil count is increased in moderate to severe preeclampsia.

REFERENCES

1. Roberts JM, Pearson G, Cutler J and Lindheimer M. Summary of the NHLBI Working Group on Research on Hypertension During Pregnancy. *Hypertension* 2003; 41:437-445.
2. Cunningham FG, Kenneth J, Steven L. Bloom et al: Hypertensive Disorders in Pregnancy. In *Williams Obstetrics*. International edn, New York: McGraw hill; 2001; chp-34, p. 567– 618.
3. Ferrazzani S, Luciano R, Garofalo S, D'Andrea V, De Carolis S, De Carolis MP, Paolucci V, Romagnoli C, Caruso A. Neonatal outcome in hypertensive disorders of pregnancy. *Early Hum Dev.* 2011; 87(6): 445-49.
4. Michelle Hladunewich, S Ananth Karumanchi, Richard Lafayette. Pathophysiology of the Clinical Manifestations of Preeclampsia. *Clin J Am Soc Nephrol* 2007; 2: 543-49.
5. Egwuatu VE. Plasma concentration of urate, urea and creatinine in Nigerian primigravidae with preeclampsia. *Trop Geogr Med* 1986; 38:11-15.
6. Mustaphi R, Gopalan S, Dhaliwal L, Sarkar AK. Hyperuricemia and pregnancy induced hypertension-reappraisal. *Indian J Med Sci* 1996, 50: 68- 71.
7. Sandhya Sivakumar, B. Vishnu Bhat and Bhawana Ashok Badhe. Effect of Pregnancy Induced Hypertension on Mothers and their babies. *Indian Journal of Pediatrics*, 2007;(74):(7):623-5.
8. Schiessl B. Inflammatory response in preeclampsia. *Mol Aspects Med* 2007; 28 (2):210-19.
9. American College of Obstetrics and Gynaecologists. Management of preeclampsia. Technical Bulletin No.1. Washington, DC; American College of Obstetrics and Gynaecology, 1986
10. Canzoneri BJ, Lewis DF, Groome L, Wang Y. Increased neutrophil numbers account for leukocytosis in women with preeclampsia. *Am J Perinatol* 2009; 26 (10):729-32.
11. Sivakumar S, Bhat BV, Badhe BA. Effect of pregnancy induced hypertension on mothers and their babies. *Indian J Pediatr* 2007; 74(7):623-25.
12. Lurie S, Frenkel E, Tuvbin Y. Comparison of the differential distribution of leukocytes in preeclampsia versus uncomplicated pregnancy. *Gynecol Obstet Invest.* 1998; 45(4):229-31.
13. Vonne J, Leif M, BERG Göran B, Jan E, Katri N, Christina K. Indications of an altered immune balance in preeclampsia: A decrease in in vitro secretion of IL-5 and IL-

10 from blood mononuclear cells and in blood basophil counts compared with normal pregnancy. Journal of

reproductive immunology 2005; 66: 69-84.

How to cite this article: Monteiro G, Subbalakshmi NK, Anupama N et. al. Total and differential leucocyte count in pregnancy induced hypertension. Int J Health Sci Res. 2013;3(12):105-109.

International Journal of Health Sciences & Research (IJHSR)

Publish your work in this journal

The International Journal of Health Sciences & Research is a multidisciplinary indexed open access double-blind peer-reviewed international journal that publishes original research articles from all areas of health sciences and allied branches. This monthly journal is characterised by rapid publication of reviews, original research and case reports across all the fields of health sciences. The details of journal are available on its official website (www.ijhsr.org).

Submit your manuscript by email: editor.ijhsr@gmail.com OR editor.ijhsr@yahoo.com